

Evaluation of end-tidal carbon dioxide role in predicting elevated SOFA scores and lactic acidosis

Daniel C. McGillicuddy · Aimee Tang ·
Lauren Cataldo · Julia Gusev · Nathan I. Shapiro

Published online: 8 January 2009
© SIMI 2009

Erratum to: Intern Emerg Med
DOI 10.1007/s11739-008-0153-z

Unfortunately, Fig. 1 of the article is mislabeled and contains erroneous data. Please find the correct figure here.

ETCO₂ and Lactate:

Decision Rule	Lactate ≥ 4	Lactate < 4	
ETCO ₂ < 35	3	53	56
ETCO ₂ ≥ 35	2	39	41
	5	92	

Sensitivity 73% (95%CI: 56-85%)
Specificity 50% (38-62%)

ETCO₂ and SOFA:

Decision Rule	SOFA ≥ 2	SOFA < 2	
ETCO ₂ < 35	25	31	56
ETCO ₂ ≥ 35	9	32	41
	34	63	

Sensitivity 73% (95%CI: 56-85%)
Specificity 50% (38-62%)

Fig. 1 This figure represents the operating characteristics for ETCO₂ at a threshold of 35 for predicting a lactate ≥ 4 mmol/L

The online version of the original article can be found under doi:[10.1007/s11739-008-0153-z](https://doi.org/10.1007/s11739-008-0153-z).

D. C. McGillicuddy (✉) · N. I. Shapiro
Department of Emergency Medicine,
Beth Israel Deaconess Medical Center,
One Deaconess Road,
WCC-2, Boston, MA 02215, USA
e-mail: dmcgilli@bidmc.harvard.edu

A. Tang · L. Cataldo
Beth Israel Deaconess Medical Center,
Boston, USA

J. Gusev
Harvard Medical School,
Boston, USA